

## A: TABLE OF BASIC DERIVATIVES

Let  $u = u(x)$  be a differentiable function of the independent variable  $x$ , that is  $u'(x)$  exists.

(A) The Power Rule :		Examples :
$\frac{d}{dx}\{u^n\} = nu^{n-1} \cdot u'$		$\frac{d}{dx}\{(x^3 + 4x + 1)^{3/4}\} = \frac{3}{4}(x^3 + 4x + 1)^{-1/4} \cdot (3x^2 + 4)$
$\frac{d}{dx}\{\sqrt{u}\} = \frac{1}{2\sqrt{u}} \cdot u'$		$\frac{d}{dx}\{\sqrt{2 - 4x^2 + 7x^5}\} = \frac{1}{2\sqrt{2 - 4x^2 + 7x^5}} (-8x + 35x^4)$
$\frac{d}{dx}\{c\} = 0$ , $c$ is a constant		$\frac{d}{dx}\{\pi^6\} = 0$ , since $\pi \cong 3.14$ is a constant.
(B) The Six Trigonometric Rules :		Examples :
$\frac{d}{dx}\{\sin(u)\} = \cos(u) \cdot u'$		$\frac{d}{dx}\{\sin(x^3)\} = \cos(x^3) \cdot 3x^2$
$\frac{d}{dx}\{\cos(u)\} = -\sin(u) \cdot u'$		$\frac{d}{dx}\{\cos(\sqrt{x})\} = -\sin(\sqrt{x}) \cdot \frac{1}{2\sqrt{x}}$
$\frac{d}{dx}\{\tan(u)\} = \sec^2(u) \cdot u'$		$\frac{d}{dx}\{\tan(\frac{5}{x^2})\} = \sec^2(5x^{-2}) \cdot (-10x^{-3})$
$\frac{d}{dx}\{\cot(u)\} = -\csc^2(u) \cdot u'$		$\frac{d}{dx}[\cot\{\sin(2x)\}] = -\csc^2\{\sin(2x)\} \cdot 2\cos(2x)$ .
$\frac{d}{dx}\{\sec(u)\} = \sec(u) \tan(u) \cdot u'$		$\frac{d}{dx}\{\sec(\sqrt[4]{x})\} = \sec(\sqrt[4]{x}) \tan(\sqrt[4]{x}) \cdot \frac{1}{4}x^{-3/4}$
$\frac{d}{dx}\{\csc(u)\} = -\csc(u) \cot(u) \cdot u'$		$\frac{d}{dx}\{\csc(8x - 7)\} = -\csc(8x - 7) \cot(8x - 7) \cdot 8$
(C) The Six Hyperbolic Rules :		Examples :
$\frac{d}{dx}\{\sinh(u)\} = \cosh(u) \cdot u'$		$\frac{d}{dx}\{\sinh(\sqrt[3]{x})\} = \cosh(\sqrt[3]{x}) \cdot \frac{1}{3}x^{-2/3}$
$\frac{d}{dx}\{\cosh(u)\} = \sinh(u) \cdot u'$		$\frac{d}{dx}\{\cosh(\sec(x))\} = \sinh\{\sec(x)\} \cdot \sec(x) \tan(x)$
$\frac{d}{dx}\{\tanh(u)\} = \operatorname{sech}^2(u) \cdot u'$		$\frac{d}{dx}[\tanh\{x^3 + \sin(x^2)\}] = \operatorname{sech}^2\{x^3 + \sin(x^2)\} \cdot (3x^2 + 2x \cos(x^2))$
$\frac{d}{dx}\{\coth(u)\} = -\operatorname{csch}^2(u) \cdot u'$		$\frac{d}{dx}\{\coth(\frac{1}{x} + 2x)\} = -\operatorname{csch}^2(\frac{1}{x} + 2x) \cdot (-\frac{1}{x^2} + 2)$
$\frac{d}{dx}\{\operatorname{sech}(u)\} = -\operatorname{sech}(u) \tanh(u) \cdot u'$		$\frac{d}{dx}\{\operatorname{sech}(9x)\} = -\operatorname{sech}(9x) \tanh(9x) \cdot 9$
$\frac{d}{dx}\{\operatorname{csch}(u)\} = -\operatorname{csch}(u) \coth(u) \cdot u'$		$\frac{d}{dx}[\operatorname{csch}\{\sinh(3x)\}] = -\operatorname{csch}\{\sinh(3x)\} \coth\{\sinh(3x)\} \cdot 3 \cosh(3x)$
(D) The Exponential & Logarithmic Rule :		Examples :
$\frac{d}{dx}\{e^u\} = e^u \cdot u'$		$\frac{d}{dx}\{e^{-x^3}\} = e^{-x^3} \cdot (-3x^2)$
$\frac{d}{dx}\{\ln u \} = \frac{u'}{u}$		$\frac{d}{dx}\{\ln x^3 + 5x + 6 \} = \frac{3x^2 + 5}{x^3 + 5x + 6}$
$\frac{d}{dx}\{a^u\} = a^u \cdot \ln(a) \cdot u' \quad , \quad a \in \mathbb{R}, a > 0, a \neq 1$		$\frac{d}{dx}\{2^{\sec(x)}\} = 2^{\sec(x)} \cdot \ln(2) \cdot \sec(x) \tan(x)$
$\frac{d}{dx}\{\log_a u \} = \frac{1}{\ln(a)} \frac{u'}{u} \quad , \quad a \in \mathbb{R}, a > 0, a \neq 1$		$\frac{d}{dx}\{\log_4 \tan(x) \} = \frac{1}{\ln(4)} \frac{\sec^2(x)}{\tan(x)}$

(E) The Six Inverse Trigonometric Functions :		Examples :
$\frac{d}{dx} \{\sin^{-1}(u)\} = \frac{u'}{\sqrt{1-u^2}}$		$\frac{d}{dx} \{\sin^{-1}(4x^2)\} = \frac{8x}{\sqrt{1-16x^4}}$
$\frac{d}{dx} \{\cos^{-1}(u)\} = -\frac{u'}{\sqrt{1-u^2}}$		$\frac{d}{dx} \{\cos^{-1}(3x)\} = -\frac{3}{\sqrt{1-9x^2}}$
$\frac{d}{dx} \{\tan^{-1}(u)\} = \frac{u'}{1+u^2}$		$\frac{d}{dx} \{\tan^{-1}(\sqrt{x})\} = \frac{1}{2\sqrt{x}} = \frac{1}{2\sqrt{x}(1+x)}$
$\frac{d}{dx} \{\cot^{-1}(u)\} = -\frac{u'}{1+u^2}$		$\frac{d}{dx} \{\cot^{-1}(e^x)\} = -\frac{e^x}{1+e^{2x}}$
$\frac{d}{dx} \{\sec^{-1}(u)\} = \frac{u'}{ u \sqrt{u^2-1}}$		$\frac{d}{dx} [\sec^{-1}(x^4)] = \frac{4x^3}{ x^4 \sqrt{x^8-1}} = \frac{4x^3}{x^4\sqrt{x^8-1}}$
$\frac{d}{dx} \{\csc^{-1}(u)\} = -\frac{u'}{ u \sqrt{u^2-1}}$		$\frac{d}{dx} \{\csc^{-1}(2x)\} = -\frac{2}{ 2x \sqrt{4x^2-1}} = -\frac{1}{ x \sqrt{4x^2-1}}$
(F) The Inverse Hyperbolic Functions :		Examples :
$\frac{d}{dx} \{\sinh^{-1}(u)\} = \frac{u'}{\sqrt{1+u^2}}$		$\frac{d}{dx} \{\sinh^{-1}(\ln(x))\} = \frac{1/x}{\sqrt{1+\ln^2(x)}}$
$\frac{d}{dx} \{\cosh^{-1}(u)\} = \frac{u'}{\sqrt{u^2-1}}$		$\frac{d}{dx} \{\cosh^{-1}(5x)\} = \frac{5}{\sqrt{25x^2-1}}$
$\frac{d}{dx} \{\tanh^{-1}(u)\} = \frac{u'}{1-u^2}$		$\frac{d}{dx} \{\tanh^{-1}(\frac{2}{x})\} = \frac{-\frac{2}{x^2}}{1-\frac{4}{x^2}} = \frac{-2}{x^2-4}$
(G) The Product and Quotient Rules :		Examples :
$\frac{d}{dx} \{uv\} = u'v + uv'$		$\frac{d}{dx} \{x^3 \ln(5x+1)\} = 3x^2 \ln(5x+1) + x^3 \frac{5}{5x+1}$
$\frac{d}{dx} \{ku\} = ku' , \quad k \text{ is a constant}$		$\frac{d}{dx} \{\frac{x^3}{4}\} = \frac{1}{4} \frac{d}{dx} \{x^3\} = \frac{1}{4} \cdot 3x^2 = \frac{3x^2}{4}$
$\frac{d}{dx} \{\frac{u}{v}\} = \frac{u'v - uv'}{v^2}$		$\frac{d}{dx} \{\frac{\tan(2x)}{x^3}\} = \frac{2 \sec^2(2x) \cdot x^3 - \tan(2x) \cdot 3x^2}{x^6}$

In table above it is assumed that  $u = u(x)$  and  $v = v(x)$  are differentiable functions

## B: TABLE OF BASIC INTEGRALS

Let  $r$ ,  $a$ ,  $b$ , and  $\beta \in \mathbb{R}$ ,  $r \neq -1$ ,  $a \neq 0$ , and  $\beta > 0$ .

<b>(A) The Power Rule :</b>	
$\int (ax + b)^r dx = \frac{(ax + b)^{r+1}}{a(r+1)} + C$	$\int x^{-5} dx = -\frac{1}{4}x^{-4} + C$ , $\int (3x - 1)^{-2} dx = \frac{(3x - 1)^{-1}}{-3} + C$
$\int dx = \int 1 dx = x + C$	$\int 7dx = 7 \int dx = 7x + C$
$\int \frac{1}{\sqrt{ax + b}} dx = \frac{2}{a} \sqrt{ax + b} + C$	$\int \frac{1}{\sqrt{x+4}} dx = 2\sqrt{x+4} + C$ .
<b>(B) The Six Trigonometric Rules :</b>	
$\int \sin(ax + b) dx = -\frac{1}{a} \cos(ax + b) + C$	$\int \sin(9x - 2) dx = -\frac{1}{9} \cos(9x - 2) + C$
$\int \cos(ax + b) dx = \frac{1}{a} \sin(ax + b) + C$	$\int \cos(3x) dx = \frac{1}{3} \sin(3x) + C$
$\int \tan(ax + b) dx = \frac{1}{a} \ln \sec(ax + b)  + C$	$\int \tan(5w - 1) dw = \frac{1}{5} \ln \sec(5w - 1)  + C$
$\int \cot(ax + b) dx = \frac{1}{a} \ln \sin(ax + b)  + C$	$\int \cot(1 - 7u) du = -\frac{1}{7} \ln \sin(1 - 7u)  + C$
$\int \sec(ax + b) dx = \frac{1}{a} \ln \sec(ax + b)  + \tan(ax + b) + C$	$\int \sec(3x) dx = \frac{1}{3} \ln \sec(3x)  + \tan(3x) + C$
$\int \csc(ax + b) dx = \frac{1}{a} \ln \csc(ax + b)  - \cot(ax + b) + C$	$\int \csc(2t) dt = \frac{1}{2} \ln \csc(2t)  - \cot(2t) + C$
<b>(C) Additional Trigonometric Rules :</b>	
$\int \sec^2(ax + b) dx = \frac{1}{a} \tan(ax + b) + C$	$\int \sec^2(2u/3) du = \frac{3}{2} \tan(2u/3) + C$
$\int \csc^2(ax + b) dx = -\frac{1}{a} \cot(ax + b) + C$	$\int \csc^2(\frac{w}{2}) dw = -\frac{1}{1/2} \cot(\frac{w}{2}) + C = -2 \cot(\frac{w}{2}) + C$
$\int \sec(ax + b) \tan(ax + b) dx = \frac{1}{a} \sec(ax + b) + C$	$\int \sec(3u) \tan(3u) du = \frac{1}{3} \sec(3u) + C$
$\int \csc(ax + b) \cot(ax + b) dx = -\frac{1}{a} \csc(ax + b) + C$	$\int \csc(5x) \cot(5x) dx = -\frac{1}{5} \csc(5x) + C$
<b>(D) The Six Hyperbolic Rules :</b>	
$\int \sinh(ax + b) dx = \frac{1}{a} \cosh(ax + b) + C$	$\int \sinh(2x - 7) dx = \frac{1}{2} \cosh(2x - 7) + C$
$\int \cosh(ax + b) dx = \frac{1}{a} \sinh(ax + b) + C$	$\int \cosh(\frac{2x}{5}) dx = \frac{5}{2} \sinh(\frac{2x}{5}) + C$
$\int \tanh(ax + b) dx = \frac{1}{a} \ln[\cosh(ax + b)] + C$	$\int \tanh(2u) du = \frac{1}{2} \ln[\cosh(2u)] + C$
$\int \coth(ax + b) dx = \frac{1}{a} \ln \sinh(ax + b)  + C$	$\int \coth(x + 3) dx = \ln \sinh(x + 3)  + C$
$\int \operatorname{sech}(ax + b) dx = \frac{2}{a} \tan^{-1}(e^{ax+b}) + C$	$\int \operatorname{sech}(3x - 6) dx = \frac{2}{3} \tan^{-1}(e^{3x-6}) + C$
$\int \operatorname{csch}(ax + b) dx = \frac{1}{a} \ln \tanh(ax + b)/2  + C$	$\int \operatorname{csch}(10t) dt = \frac{1}{10} \ln \tanh(5t)  + C$

(E) Additional Hyperbolic Rules :	Examples
$\int \operatorname{sech}^2(ax+b)dx = \frac{1}{a} \tanh(ax+b) + C$	$\int \operatorname{sech}^2(4w)dw = \frac{1}{4} \tanh(4w) + C$
$\int \operatorname{csch}^2(ax+b)dx = -\frac{1}{a} \coth(ax+b) + C$	$\int \operatorname{csch}^2(2u)du = -\frac{1}{2} \coth(2u) + C$
$\int \operatorname{sech}(ax+b)\tanh(ax+b)dx = -\frac{1}{a} \operatorname{sech}(ax+b) + C$	$\int \operatorname{sech}(3x)\tanh(3x)dx = -\frac{\operatorname{sech}(3x)}{3} + C$
$\int \operatorname{csch}(ax+b)\coth(ax+b)dx = -\frac{1}{a} \operatorname{csch}(ax+b) + C$	$\int \operatorname{csch}(\frac{x}{3})\coth(\frac{x}{3})dx = -3\operatorname{csch}(x/3) + C$
(F) Exponential /Logarithmic Rules :	Examples :
$\int e^{ax+b}dx = \frac{1}{a}e^{ax+b} + C$	$\int e^{7x}dx = \frac{1}{7}e^{7x} + C$
$\int k^{ax+\beta}dx = \frac{1}{a\ln(k)}.k^{ax+\beta} + C , 0 < k \in \mathbb{R}, k \neq 1.$	$\int 2^{10x-17}dx = \frac{1}{10\ln 2} 2^{10x-17} + C$
$\int \frac{1}{ax+b}dx = \frac{1}{a} \ln ax+b  + C$	$\int \frac{1}{2x-3}dx = \frac{1}{2} \ln 2x-3  + C$
(G) The Three Inverse Trigonometric Functions :	Examples :
$\int \frac{1}{\sqrt{\beta^2-x^2}}dx = \sin^{-1}(\frac{x}{\beta}) + C$	$\int \frac{1}{\sqrt{16-x^2}}dx = \sin^{-1}(x/4) + C$
$\int \frac{1}{\beta^2+x^2}dx = \frac{1}{\beta} \tan^{-1}(\frac{x}{\beta}) + C$	$\int \frac{1}{3+x^2}dx = \frac{1}{\sqrt{3}} \tan^{-1}(\frac{x}{\sqrt{3}}) + C$
$\int \frac{1}{x\sqrt{x^2-\beta^2}}dx = \frac{1}{\beta} \sec^{-1}(\frac{x}{\beta}) + C , x > \beta$	$\int \frac{1}{x\sqrt{x^2-4}}dx = \frac{1}{2} \sec^{-1}(\frac{x}{2}) + C , x > 2.$
(H) The Three Inverse Hyperbolic Functions :	Examples :
$\int \frac{1}{\sqrt{\beta^2+x^2}}dx = \sinh^{-1}(\frac{x}{\beta}) + C$	$\int \frac{1}{\sqrt{1+x^2}}dx = \sinh^{-1}(x) + C$
$\int \frac{1}{\sqrt{x^2-\beta^2}}dx = \cosh^{-1}(\frac{x}{\beta}) + C$	$\int \frac{1}{\sqrt{x^2-5}}dx = \cosh^{-1}(x/\sqrt{5}) + C$
$\int \frac{1}{\beta^2-x^2}dx = \frac{1}{\beta} \tanh^{-1}(\frac{x}{\beta}) + C ,  x  < \beta$	$\int \frac{1}{36-x^2}dx = \frac{1}{6} \tanh^{-1}(\frac{x}{6}) + C ,  x  < 6$
(I) The Fundamental Theorems	Examples :
$\int_a^b f(x)dx = g(x) _{x=a}^{x=b} = g(b) - g(a)$	$\int_e^{e^3} \frac{1}{x}dx = \ln x  _{x=e}^{x=e^3} = \ln(e^3) - \ln(e) = 3 - 1 = 2$
$\frac{d}{dx} \left\{ \int_{u(x)}^{v(x)} F(t) dt \right\} = F(v(x)).v'(x) - F(u(x)).u'(x)$	$\frac{d}{dx} \left\{ \int_x^{x^2} \cos(t^2)dt \right\} = \cos(x^4).2x - \cos(x^2).1$

In table above it is assumed that :

(1) The function  $f(x)$  is continuous on  $[a, b]$  and  $\int f(x) dx = g(x) + C$ .

(2) The functions  $u(x)$  and  $v(x)$  are differentiable and  $\int_{u(x)}^{v(x)} F(t) dt$  exists.

## C: BASIC TRIGONOMETRIC IDENTITIES

### GROUP (A) :

$$(i) \tan(\theta) = \frac{\sin(\theta)}{\cos(\theta)} \quad (ii) \cot(\theta) = \frac{\cos(\theta)}{\sin(\theta)}$$

$$(iii) \sec(\theta) = \frac{1}{\cos(\theta)} \quad (iv) \csc(\theta) = \frac{1}{\sin(\theta)}$$

### GROUP (B) :

$$(i) \cos^2(\theta) + \sin^2(\theta) = 1 \quad (ii) 1 + \tan^2(\theta) = \sec^2(\theta) \quad (iii) \cot^2(\theta) + 1 = \csc^2(\theta)$$

### GROUP (C) :

$$(i) \sin(2\theta) = 2 \sin(\theta) \cos(\theta) \quad (ii) \cos(2\theta) = 2 \cos^2(\theta) - 1 \quad (iii) \cos(2\theta) = 1 - 2 \sin^2(\theta)$$

$$(iv) \cos^2(\theta) = \frac{1}{2}[1 + \cos(2\theta)] \quad (v) \sin^2(\theta) = \frac{1}{2}[1 - \cos(2\theta)]$$

### GROUP (D)

$$(i) \sin(-\theta) = -\sin(\theta) \quad (ii) \cos(-\theta) = \cos(\theta) \quad (iii) \tan(-\theta) = -\tan(\theta).$$

### GROUP(E)

$$(i) \cos(\theta \pm \phi) = \cos(\theta) \cos(\phi) \mp \sin(\theta) \sin(\phi)$$

$$(ii) \sin(\theta \pm \phi) = \sin(\theta) \cos(\phi) \pm \cos(\theta) \sin(\phi)$$

### GROUP (F)

$$(i) \cos(\theta) \cos(\phi) = \frac{1}{2}[\cos(\theta - \phi) + \cos(\theta + \phi)]$$

$$(ii) \sin(\theta) \sin(\phi) = \frac{1}{2}[\cos(\theta - \phi) - \cos(\theta + \phi)]$$

$$(iii) \sin(\theta) \cos(\phi) = \frac{1}{2}[\sin(\theta - \phi) + \sin(\theta + \phi)]$$

## D: SPECIAL TRIGONOMETRIC EQUATIONS

$$(i) \sin(x) = 0 \Rightarrow x = n\pi \quad (ii) \cos(x) = 0 \Rightarrow x = \frac{(2n-1)}{2}\pi$$

where  $n$  is an integer :  $n = 0, \pm 1, \pm 2, \pm 3, \dots$

## E: HYPERBOLIC FUNCTIONS

$$(i) \sinh(x) = \frac{1}{2}[e^x - e^{-x}] \quad (ii) \cosh(x) = \frac{1}{2}[e^x + e^{-x}] \quad (iii) \tanh(x) = \frac{\sinh(x)}{\cosh(x)}$$

$$(iv) \coth(x) = \frac{\cosh(x)}{\sinh(x)} \quad (v) \operatorname{sech}(x) = \frac{1}{\cosh(x)} \quad (vi) \operatorname{csch}(x) = \frac{1}{\sinh(x)}$$

$$(vii) \cosh^2(x) - \sinh^2(x) = 1 \quad (viii) 1 - \tanh^2(x) = \operatorname{sech}^2(x) \quad (ix) \coth^2(x) - 1 = \operatorname{csch}^2(x)$$

## F: PROPERTIES OF LOGARITHMS

Let  $x$  and  $y$  be positive real numbers.

$$(i) \ln(x) + \ln(y) = \ln(xy) \quad (ii) \ln(x) - \ln(y) = \ln(\frac{x}{y}) \quad (iii) \ln(x^m) = m \ln(x).$$

$$(iv) \ln(e^k) = k \quad (v) e^{\ln(x)} = x \quad (vi) \ln(1) = 0, \ln(e) = 1.$$

## G: SPECIAL VALUES

$$(i) \sin(0) = 0 \quad (ii) \cos(0) = 1 \quad (iii) \tan(0) = 0$$

$$(iv) \sinh(0) = 0 \quad (v) \cosh(0) = 1 \quad (vi) \tanh(0) = 0$$

(vii)  $\sin(n\pi) = 0$  and  $\cos(n\pi) = (-1)^n$ , provided that "  $n$  " is an integer.

END