

CO₂ as a primary driver of Phanerozoic climate: COMMENT

Nir Shaviv

Racah Institute of Physics, Hebrew University of Jerusalem, Jerusalem, 91904 Israel

Jan Veizer

Institut für Geologie, Mineralogie und Geophysik, Ruhr-Universität Bochum, 44801 Bochum, Germany, and Department of Earth Sciences, University of Ottawa, Ottawa, ON K1N 6N5 Canada

Royer et al. (2004) introduce a seawater pH correction to the Phanerozoic temperature reconstruction based on $\delta^{18}\text{O}$ variations in marine fossils. Although this correction is a novel idea and it is likely to have played some role in offsetting the $\delta^{18}\text{O}$ record, we show that (a) the correction cannot be as large as claimed by Royer et al., (b) irrespective of the size of the correction, a CO₂ signature cannot possibly be seen in the data, and (c) even though the CO₂ signature cannot be seen, the pH correction implies only a somewhat higher global temperature sensitivity than that in Shaviv and Veizer (2003), a sensitivity that is consistent with a “black body Earth,” but only marginally with the lower limit of the Intergovernmental Panel on Climate Change (IPCC) range.

INTRODUCTION

Royer et al. (2004) citing the standard references from the IPCC (2001) volume assert that CO₂ is the principal climate driver on modern as well as geological time scales. Although some of these cited publications (e.g., Mann et al., 1998) were questioned recently (Soon and Baliunas, 2003; McIntire and McKittrick, 2003), it is not our intention to scrutinize their methodology. We only would like to point out that these publications implicitly assume that the various radiative forcings at play do not include an amplifier to solar activity. However, a large body of empirical evidence, published mostly subsequent to the IPCC summary (e.g., Bond et al., 2001; Neff et al., 2001; Solanki, 2002; Rind, 2002; Foukal, 2002; Usoskin et al., 2003) suggests that such an amplifier exists, and it is most likely in the form of solar modulation of the cosmic ray flux (CRF) (e.g., Tinsley and Deen, 1991; Svensmark, 1998, 2000; Marsh and Svensmark, 2000, 2003; Palle Bago and Butler, 2000; Egorova et al., 2000). Note that Royer et al. do not dispute the existence of the CRF/temperature correlation of Shaviv and Veizer (2003), only its role relative to that of CO₂. The CRF effect should therefore be incorporated as a complementary factor into the existing climate models.

Once this solar amplification is included, the paleoclimate data is consistent with a solar (direct and indirect) contribution of 0.32 ± 0.11 °C toward global warming over the past century. While these results imply different conclusions than the IPCC scenarios, they are not necessarily contradictory given the differences in basic assumptions. Note also that when the temperature trend over the past century is compared to the monotonic increase in greenhouse gases versus the non-monotonic increase in solar activity, the best fit suggests that more than half of the warming should be attributed to solar activity and somewhat less to greenhouse gases (GHG) (Soon et al., 1996). A similar comparison with the Total Solar Irradiance (TSI) flux led Foukal (2002) to state that the TSI can explain 80% of the 11-year smoothed variance in the centennial global temperature trend. Such empirical observations strongly suggest that celestial forcing played an important role in the climate history of our planet.

Royer et al. (2004) cite the unquestionably impressive correlation of CO₂ and polar temperatures in Antarctic ice cores (Barnola et al., 1987; Petit et al., 1999) as a proof that CO₂ is the climate driver on $\sim 10^5$ year time scales. This correlation, however, cannot distinguish by itself between cause and effect. More important, recent high-resolution data (Mudelsee, 2001; Clarke, 2003) demonstrates that the rises in CO₂ lag by centuries (i.e., much smaller than the 10^5 year time scale) behind the increases in

temperature. Clearly, the CO₂ may act as a temperature amplifier, but not as the driver for climate changes that happened centuries earlier.

On still longer, Phanerozoic time scales, Royer et al. (2004) cite the “close correspondence between CO₂ and temperature” as advocated in Crowley and Berner (2001). A simple visual inspection of the alternatives (Fig. 1) shows that it is the “four-hump” reconstruction of $\delta^{18}\text{O}$ based paleotemperatures (Veizer et al., 2000) and the CRF (Shaviv and Veizer, 2003) that correlate closely with the paleoclimate history based on sedimentary indicators (<http://www.scotese.com/climate.htm>; Boucot and Gray, 2001), and not the “two-hump” GEOCARB III reconstruction (Royer et al., 2004) of atmospheric CO₂. Note also that the GEOCARB III results in high atmospheric CO₂ levels for most of the Paleozoic and mid-Mesozoic. In order to explain the recurring cold intervals during these times, one has to resort to a multitude of special pleadings. The late Ordovician glaciation at an apparent ~ 5000 ppm CO₂ is a classic example.

Royer et al. (2004) discount the validity of all other Phanerozoic CO₂ reconstructions, citing the general agreement of the GEOCARB III trend with estimates based on CO₂ proxies. Considering the many assumptions and large uncertainties in the proxy data (Royer et al., 2001) this statement may need future confirmation, but we are prepared to accept it at this stage. Nevertheless, we (Shaviv and Veizer, 2003) believe that in view of the above, it is necessary to test all presently available Phanerozoic CO₂ scenarios. The outcome, however, is more or less the same for all alternatives—a lack of correlation with the $\delta^{18}\text{O}$ based paleotemperatures as well as with the paleoclimate trend based on sediments.

SPECIFIC MODELING COMMENTS CONCERNING THE pH CORRECTION OF ROYER ET AL.

The pH effect introduced by Royer et al. (2004) may well have played a role in distorting the $\delta^{18}\text{O}$ data, such that a reconstructed temperature should take it into consideration. We show here, however, that while the inferred influence of CO₂ (and global temperature sensitivity in general) is somewhat higher once the pH effect is considered, the qualitative conclusions reached in Shaviv and Veizer (2003) are generally the same. This is because of several critical points elaborated here.

THEORETICAL BACKGROUND

Zeebe (1999) considered the possible bias of $\delta^{18}\text{O}$ -derived temperature due to the carbonate ion effect on oxygen isotope incorporation in carbonates controlled by temporal changes in the pH. Zeebe (2001) derived the expression $\Delta T_{\text{pH}} = bs\Delta(\text{pH})$ for the correction of the reconstructed temperature using $\delta^{18}\text{O}$ of carbonates. s is the linear coefficient between $\delta^{18}\text{O}$ and pH from theory and experiments ($s = -1.42\%$ per unit pH). b is the linear coefficient between $\delta^{18}\text{O}$ and temperature. The value of b taken by Royer et al. (2004) is -4.90 °C per 1‰ of $\delta^{18}\text{O}$. However, this does not allow for an ice-volume effect. If one assumes no ice-caps at the peak of the greenhouse epochs, and ice-sheets twice as large as today at the height of the icehouses, then as much as 2‰ of the 3‰–5‰ range in the $\delta^{18}\text{O}$ data should be attributed to an ice-volume effect and not to temperature (Guilderson et al. 1994; Savin, 1977; Veizer et al., 2000). Thus, on average over the Phanerozoic, a more appropriate value is b of -2.0 °C per 1‰ of $\delta^{18}\text{O}$.

Once this is considered, the pH correction to the $\delta^{18}\text{O}$ -reconstructed temperature is

$$\Delta T_{\text{pH}} = a \{ \log R_{\text{CO}_2} + \log \Lambda(t) - \log \Omega(t) \}, \quad (1)$$

where R_{CO_2} is the atmospheric partial pressure of CO₂ relative to today, $\Lambda(t)$ is $(\text{Ca}(t)/(\text{Ca})(0))$ —the mean concentration of dissolved calcium in the water relative to today, while $\Omega(t)$ is $[\text{Ca}^{++}][\text{CO}_3]/K_{\text{sp}}$ at time t relative to today. The value of a obtained in Royer et al. (2004) assumes no ice-volume correction, yielding $a = 3.4$ °C. If we take the smaller b , given the likely ice-volume effect, we obtain $a = 1.4$ °C. This correction is important, as documented below.

Methodology

Royer et al. (2004) apply the pH correction to obtain a corrected reconstructed temperature of:

$$\Delta T_{O_{18}} = b\delta^{18}O + \Delta T_{pH} = b\delta^{18}O + a \{ \log R_{CO_2} + \log \Lambda(t) - \log \Omega(t) \}. \quad (2)$$

When comparing to the variable CO₂ and cosmic ray flux (CRF), one essentially compares the temperature to a modeled temperature of the form:

$$\Delta T_{model} = A + Bt + C \log_2 R_{CO_2} + Dg(\Phi_{CRF}(t)), \quad (3)$$

where the last term is the contribution from the CRF as described in Shaviv and Veizer (2003). The first two terms allow for systematic variations in temperature, some of which are poorly known (e.g., secular solar luminosity increase, systematic error in the $\delta^{18}O$ detrending, etc.).

The comparison of Royer et al. (2004) is basically between $\Delta T_{O_{18}}$ and ΔT_{model} . The $\Lambda(t)$ and $\Omega(t)$ terms are relatively minor. Thus, the predominant pH correction is from the R_{CO_2} term. Without this term, Shaviv and Veizer (2003) find that there is no apparent correlation between $\Delta T_{O_{18}}$ and ΔT_{model} . In other words, the correlation (i.e., agreement) between the GEOCARB III reconstruction and pH corrected temperature was obtained through bootstrapping. Any R_{CO_2} reconstruction would have contributed to a ΔT_{pH} correction, making the $\Delta T_{O_{18}}$ correlate with the R_{CO_2} . That is to say, the temperature reconstruction cannot be used to claim that a R_{CO_2} fingerprint is seen. In fact, any R_{CO_2} used will imply a correlation between CO₂ and ΔT of about $C \sim a \log_{10} 2 = 0.4$ °C (or 1.0 °C if no ice-volume effect is present), even for the extreme case where absolutely no correlation exists between $\delta^{18}O$ and R_{CO_2} . (In reality, of course, the statement that there is “no apparent correlation” between $\delta^{18}O$ and R_{CO_2} is statistically limited, such that C has errors around the above value.)

Because the correction to $\Delta T_{O_{18}}$ includes R_{CO_2} , a more appropriate approach should be as follows.

Let us define an “uncorrected” temperature as

$$\Delta T' \equiv \Delta T - a \log R_{CO_2}. \quad (4)$$

This uncorrected temperature does not depend on R_{CO_2} for its reconstruction

$$\Delta T'_{O_{18}} = b\delta^{18}O + \Delta T_{pH} - a \log R_{CO_2} = b\delta^{18}O + a \{ \log \Lambda(t) - \log \Omega(t) \}. \quad (5)$$

On the other hand, when we model this uncorrected temperature, we should do so with

$$\Delta T'_{model} = A + Bt + (C - a \log_{10} 2) \log_2 R_{CO_2} + Dg(\Phi_{CRF}(t)). \quad (6)$$

We added the $(a \log_{10} R_{CO_2}) = (a \log_{10} 2) \log_2 R_{CO_2}$ term such that we will keep the meaning of C , which is the tropical temperature increase associated with a doubled R_{CO_2} . In other words, the appropriate method to perform this analysis is to repeat the analysis of Shaviv and Veizer (2003), while correcting for the relatively small $\log \Lambda(t)$ and $\log \Omega(t)$ terms. To first approximation, if the latter small terms are neglected, the values of “ C ” in Shaviv and Veizer (2003) should simply be replaced with $(C - a \log_{10} 2)$. Thus, we should add the value $a \log_{10} 2 = 0.4$ °C to the results of Shaviv and Veizer (2003), or ~ 0.6 °C if we wish to correct the estimates for ΔT_{x_2} , the globally averaged temperature increase associated with a doubled R_{CO_2} .

The lack of a correlation between $\delta^{18}O$ and R_{CO_2} originates from the fact that $(C - a \log_{10} 2)$ happens to be coincidentally close to 0. In other words, the pH correction to $\delta^{18}O$ and ΔT happens to be similar to the tropical temperature sensitivity to changes in R_{CO_2} (without the pH correction, the preferred value for C in the absence of correlation is not $a \log_{10} 2$, but 0). Scientifically, this is somewhat unfortunate, because without this coincidence, the $\delta^{18}O$ signal would have had a clear correlation with the R_{CO_2} signal, and the R_{CO_2} fingerprint would have been discernible. Note

nevertheless that for a global temperature sensitivity much higher than discussed above, the CO₂ signal in the $\delta^{18}O$ data should have been present, but this is not the case.

If we repeat the analysis of Shaviv and Veizer (2003), and consider also the effects of $a \{ \log \Lambda(t) - \log \Omega(t) \}$, such correction introduces a small positive correlation between $\Delta T'_{O_{18}}$ and $\Delta T'_{model}$.

We obtain:

$C = 0.69$ °C (or an upper limit of 1.12, 1.42, and 1.73 °C at 68%, 90%, and 99% confidence levels, and a lower limit of 0.39, 0.10, and -0.21 °C, respectively).

Here, the CRF can explain 70% of the variance in ΔT (additional 11% is due to measurement error in $\delta^{18}O$). That is, the CRF is still the main climate driver, while CO₂ plays only a secondary role.

If we ignore the ice-volume effect and take a large a , we obtain:

$C = 1.29$ °C (or an upper limit of 1.64, 1.95, and 2.26 °C at 68%, 90%, and 99% confidence levels, and a lower limit of 1.07, 0.77, and 0.45 °C, respectively).

In this case, the variance in ΔT is larger, but so is the contribution of CO₂. One obtains that either the CRF or the CO₂ alone can explain 49% of the total variance, while together, they can explain 83% of the variance. In this case, the relative role of CO₂ is as high as that of the CRF, but the pH correction is unrealistically high, due to the neglected ice-volume correction.

The above values for C are somewhat larger than those in Shaviv and Veizer (2003); nevertheless, even with the neglected ice-volume effect, the value of C obtained is still relatively small. The best estimate for the global temperature sensitivity with ice-volume effect is $\Delta T_{x_2} \sim 1.1$ °C for $C = 0.69$ °C, and without it ~ 1.9 °C.

CONCLUSIONS

The analysis of Royer et al. (2004) assumes an unrealistically high pH correction. First, it neglects the ice-volume effect, which changes the relation between $\delta^{18}O$ and ΔT . Second, this large pH correction implies high temperatures for seawater even during times of extensive glaciations.

Moreover, the analysis of Royer et al. (2004) consists of bootstrapping, by introducing a correction to ΔT that is an implicit function of R_{CO_2} . It is then not surprising that a correlation between ΔT and R_{CO_2} is obtained. This would be the case irrespective of the R_{CO_2} model utilized.

A proper analysis, which avoids this bootstrapping and considers a more realistic pH correction, shows that the global temperature sensitivity to CO₂ is still relatively small. In summary, while we acknowledge that the proposition of Royer et al. (2004) has some merit and likely will result in some modification of the $\delta^{18}O$ signal, the CRF still remains the primary climate driver for any realistic pH correction (see also Wallmann, 2004). Even for the scenario that entirely disregards the ice-volume effect, the impact of CRF would still be at par with that of CO₂.

REFERENCES CITED

- Barnola, J.M., Raynaud, D., Korotkevich, Y.S., and Lorius, C., 1987, Vostok ice core provides 160,000-year record of atmospheric CO₂; *Nature*, v. 329, p. 408–414, doi: 10.1038/329408A0.
- Bond, G., Kromer, B., Beer, J., Muscheler, R., Evans, M.N., Showers, W., Hoffmann, S., Lotti-Bond, R., Hajdas, I., and Bonani, G., 2001, Persistent solar influence on North Atlantic climate during the Holocene; *Science*, v. 294, p. 2130–2136, doi: 10.1126/SCIENCE.1065680.
- Boucot, A.J., and Gray, J., 2001, A critique of Phanerozoic climatic modes involving changes in the CO₂ content of the atmosphere; *Earth Science Reviews*, v. 56, p. 1–159, doi: 10.1016/S0012-8252(01)00066-6.
- Clarke, T., 2003, Bubbles prompt climate-change rethink; *Nature*, *Science Update* (<http://www.nature.com/nsu/pdf/030310/030310-12.html>).
- Crowley, T.J., and Berner, R.A., 2001, CO₂ and climate change; *Science*, v. 292, p. 870–872, doi: 10.1126/SCIENCE.1061664.




Figure 1. The temporal reconstruction of (A) Phanerozoic climate based on the paleogeographic distribution of climate sensitive sediments and fossils (<http://www.scotese.com/climate.htm>; Boucot and Gray, 2001), (B) the trend in cosmic ray flux and paleotemperatures based on $\delta^{18}\text{O}$ (Shaviv and Veizer, 2003), and (C) the reconstructions of global temperatures and atmospheric CO_2 from GEOCARB models and proxies (Crowley and Berner, 2001). Note that the “5-hump” pattern in A is due to the 600 Ma time span, as opposed to only the 500 Ma in B. This is because Scotese incorporates sedimentation records from before the Cambrian, while the $\delta^{18}\text{O}$ reconstruction requires skeletal fossils that did not exist prior to the Cambrian explosion of life forms.

Egorova, L.Y., Vovk, V.Y., and Troshichev, O.A., 2000, Influence of variations of the cosmic rays on atmospheric pressure and temperature in the southern geomagnetic pole region: *Journal of Atmospheric Solar-Terrestrial Physics*, v. 62, p. 955, doi: 10.1016/S1364-6826(00)00080-8.

Foukal, P., 2002, A comparison of variable solar total and ultraviolet irradiance outputs in the 20th century: *Geophysical Research Letters*, v. 29, p. 411–414, 2089, doi: 10.2912002GL015474.

Guilderson, T.P., Fairbanks, R.G., and Rubenstone, J.L., 1994, Tropical temperature variations since 20,000 years ago: modulating interhemispheric climate change: *Science*, v. 263, p. 663–665.

Intergovernmental Panel on Climate Change (IPCC), 2001, Contribution of Working Group I to the third assessment report of the IPCC, Houghton, J.T., et al., eds., *Climate change 2001: The scientific basis*: Cambridge, Cambridge University Press, 892 p.

Mann, M.E., Bradley, R.S., and Hughes, M.K., 1998, Global-scale temperature patterns and climate forcing over the past six centuries: *Nature*, v. 392, p. 779–787, doi: 10.1038/33859.

Marsh, N., and Svensmark, H., 2000, Cosmic rays, clouds, and climate: *Space Science Reviews*, v. 94, p. 215–230, doi: 10.1023/A:1026723423896.

Marsh, N., and Svensmark, H., 2003, Galactic cosmic ray and El Niño southern oscillation trends in International Satellite Cloud Climatology Project D2 low-cloud properties: *Journal of Geophysical Research, Atmospheres*, v. 108, no. D6, p. 4195, doi: 10.1029/2001JD001264.

McIntire, S., and McKittrick, R., 2003, Corrections to the Mann et al. (1998) proxy database and northern hemispheric average temperature series: *Energy & Environment*, v. 14, p. 751–771.

Mudelsee, M., 2001, The phase relations among atmospheric CO_2 content, temperature and global ice volume over the past 420 ka: *Quaternary Science Reviews*, v. 20, p. 583–589, doi: 10.1016/S0277-3791(00)00167-0.

Neff, U., Burns, S.J., Mangini, A., Mudelsee, M., Fleitmann, D., and Matter, A., 2001, Strong coherence between solar variability and the monsoon in Oman between 9 and 6 kyr ago: *Nature*, v. 411, p. 290–293, doi: 10.1038/35077048.

Palle Bago, E., and Butler, J., 2000, The influence of cosmic terrestrial clouds and global warming: *Astronomy & Geophysics*, v. 41, p. 18–22.

Petit, J.R., Jouzel, J., Raynaud, D., Barkov, N.I., Barnola, J.-M., Basile, I., Benders, M., Chappellaz, J., Davis, M., Delayque, G., Delmotte, M., Kotlyakov, V.M., Legrand, M., Lipenkov, V.Y., Lorius, C., Pépin, L., Ritz, C., Saltzman, E., and Stevenard, M., 1999, Climate and atmospheric history of the past 420,000 years from the Vostok ice core, Antarctica: *Nature*, v. 399, p. 429–436, doi: 10.1038/20859.

Rind, D., 2002, The sun’s role in climate variations: *Science*, v. 296, p. 673–677, doi: 10.1126/SCIENCE.1069562.

Royer, D.L., Berner, R.A., Montañez, I.P., Tabor, N.J., and Beerling, D.J., 2004, CO_2 as a primary driver of Phanerozoic climate: *GSA Today*, v. 14, no. 3, p. 4–10, doi: 10.1130/1052-5173(2004)014<0004:CAAPDO>2.0.CO;2.

Savin, S.M., 1977, The history of the earth’s surface temperature during the past 100 million years: *Annual Reviews of Earth and Planetary Science*, v. 5, p. 319–355, doi: 10.1146/ANNUREV.EA.05.050177.001535.

Shaviv, N.J., and Veizer, J., 2003, Celestial driver of Phanerozoic climate?: *GSA Today*, v. 13, no. 7, p. 4–10, doi: 10.1130/1052-5173(2003)0132.0.CO;2.

Solanki, S.K., 2002, Solar variability and climate change: is there a link?: *Astronomy and Geophysics*, v. 43, p. 5.9–5.13.

- Soon, W.H., and Baliunas, S.L., 2003, Proxy climate and environmental changes of the past 100 years: *Climate Research*, v. 23, p. 89–110.
- Soon, W.H., Posmentier, E.S., and Baliunas, S.L., 1996, Inference of solar irradiance variability from terrestrial temperature changes, 1880–1993: An astrophysical application of the sun-climate connection: *The Astrophysical Journal*, v. 472, p. 891, doi: 10.1086/178119.
- Svensmark, H., 1998, Influence of cosmic rays on earth's climate: *Physical Review Letters*, v. 81, p. 5027–5030, doi: 10.1103/PHYSREVLETT.81.5027.
- Svensmark, H., 2000, Cosmic rays and earth's climate: *Space Science Reviews*, v. 93, p. 175–185, doi: 10.1023/A:1026592411634.
- Tinsley, B.A., and Deen, G.W., 1991, Apparent tropospheric response to MeV-GeV particle flux variations: A connection via electrofreezing of supercooled water in high-level clouds?: *Journal of Geophysical Research*, v. 96, p. 22,283–22,296.
- Usoskin, I.G., Solanki, S.K., Schüssler, M., Mursula, K., and Alanko, K., 2003, A millennium scale sunspot number reconstruction: evidence for an unusually active sun since the 1940's: *Physical Review Letters*, v. 91, p. 211101-1–211101-4.
- Veizer, J., Godderis, Y., and François, L.M., 2000, Evidence for decoupling of atmospheric CO₂ and global climate during the Phanerozoic eon: *Nature*, v. 408, p. 698–701, doi: 10.1038/35047044.
- Wallmann, K., 2004, Impact of atmospheric CO₂ and galactic cosmic radiation on Phanerozoic climate change and the marine δ¹⁸O record: *Geochemistry, Geophysics, Geosystems* (in press).
- Zeebe, R.E., 1999, An explanation of the effect of seawater carbonate concentration on foraminiferal oxygen isotopes: *Geochimica et Cosmochimica Acta*, v. 63, p. 2001–2007, doi: 10.1016/S0016-7037(99)00091-5.
- Zeebe, R.E., 2001, Seawater pH and Isotopic paleotemperatures of Cretaceous Oceans: *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 170, p. 49–57, doi: 10.1016/S0031-0182(01)00226-7.

MANUSCRIPT SUBMITTED FEBRUARY 17, 2004; ACCEPTED APRIL 27, 2004.