

Comment and Reply

CO₂ as a primary driver of Phanerozoic climate

Dana L. Royer, Robert A. Berner, Isabel P. Montañez, Neil J. Tabor, and David J. Beerling, *GSA Today*, v. 14, no. 3, p. 4–10.

Comment

Nir Shaviv, *Racah Institute of Physics, Hebrew University of Jerusalem, Jerusalem, 91904 Israel*

Ján Veizer, *Institut für Geologie, Mineralogie und Geophysik, Ruhr-Universität Bochum, 44801 Bochum, Germany, and Ottawa-Carleton Geoscience Centre, University of Ottawa, Ottawa, Ontario, K1N 6N5 Canada*

Royer et al. (2004), in their recent article in *GSA Today*, dispute the conclusion of Shaviv and Veizer (2003) that celestial forcing may have been the major driver of Phanerozoic climate, arguing instead for CO₂ as the dominant force. The δ¹⁸O of calcareous shells reflects the ambient temperature of seawater and the quantity of water locked in the polar ice caps, each contributing about one half to this signal, but lately it has been realized that seawater pH drives the δ¹⁸O in the opposite direction. Royer et al. utilized this for reconciling the GEOCARB III and the δ¹⁸O trend of Veizer et al. (2000) by assuming that any discrepancy of the two variables is due to pH. This is an interesting proposition that has some merit, but note that such a correction is entirely arbitrary, because we do not have any constraints for the pH of Phanerozoic seawater, except possibly some boron isotopes for the youngest portion of this record. Moreover, the pH correction itself basically reflects the GEOCARB III CO₂ model, such that the correlation obtained between the corrected δ¹⁸O and CO₂ cannot be claimed to be a CO₂ fingerprint. Furthermore, the model of Royer et al. does not consider the mitigating “ice volume” effect. Once included, the required pH correction (and GEOCARB III CO₂ levels) would have to be about doubled for CO₂ climate “driving” to be on par with the CRF. For all these, and the reasons listed in our detailed response (see www.gsjournals.org; go to “Online Journals,” then “Online Forum”), we argue that the δ¹⁸O trend is still chiefly a reflection of the temperature history of the past oceans, controlled principally by the celestial driver.

REFERENCES CITED

Royer, D.L., Berner, R.A., Montañez, I.P., Tabor, N.J., and Beerling, D.J., 2004, CO₂ as a primary driver of Phanerozoic climate: *GSA Today*, v. 14, no. 3, p. 4–10.

Shaviv, N.J., and Veizer, J., 2003, Celestial driver of Phanerozoic climate?: *GSA Today*, v. 13, no. 7, p. 4–10.

Veizer, J., Godderis, Y., and François, L.M., 2000, Evidence for decoupling of atmospheric CO₂ and global climate during the Phanerozoic eon: *Nature*, v. 404, p. 698–701.

Online Comments and Replies

Visit www.gsjournals.org for comments and replies on *GSA Today* articles. Click on “Online Journals” then “Online Forum,” then “GSA Today: Comments and Replies.”

Reply

Dana L. Royer, *Department of Geosciences and Institutes of the Environment, Pennsylvania State University, University Park, Pennsylvania 16802, USA, droyer@psu.edu*

Robert A. Berner, *Department of Geology and Geophysics, Yale University, New Haven, Connecticut 06520, USA*

Isabel P. Montañez, *Department of Geology, University of California, Davis, California 95616, USA*

Neil J. Tabor, *Department of Geological Sciences, Southern Methodist University, Dallas, Texas 75275, USA*

David J. Beerling, *Department of Animal and Plant Sciences, University of Sheffield, Sheffield S10 2TN, UK*

A major purpose of our paper was to show that paleotemperature records do not support the presence of prolonged cold intervals (similar in magnitude and duration to the Permo-Carboniferous glaciation) during the Ordovician/Silurian (460–400 Ma) and mid-Mesozoic (220–120 Ma), as required by the δ¹⁸O record of Veizer et al. (1999). The fact that our pH-corrected curve better matches the more robust record of glacial sediments should be taken as a positive result, considering an alternative explanation of diagenesis. Crucially, diagenetic overprinting can be present even when screening samples using the petrographic and trace element criteria adopted by Veizer et al. (1999) (cf. Montañez et al., 2000; Wenzel, 2000; Mii et al., 2001; Pearson et al., 2001).

Shaviv and Veizer claim that we correlate the CO₂ and pH-corrected temperature records. This is simply untrue. Also, their “ice volume” effect, to be applied to our pH correction, is based on an average slope for the change in temperature with δ¹⁸O, for the entire Phanerozoic. This approach is incorrect. The ice volume correction is applicable only to periods of major glaciation, but at these times our pH correction was minor.

The correspondence between the Phanerozoic records of atmospheric CO₂ and glacial sediments, and the revision of the δ¹⁸O paleotemperature record toward values better matching the glacial sediment record, strongly implicate CO₂ as a primary driver of climate over these timescales. Cosmic ray flux is likely only of second-order significance (see also Rahmstorf et al., 2004).

REFERENCES CITED

Mii, H.-S., Grossman, E.L., Yancey, T.E., Chuvashov, B., and Egorov, A., 2001, Isotopic records of brachiopod shells from the Russian Platform—evidence for the onset of mid-Carboniferous glaciation: *Chemical Geology*, v. 175, p. 133–147.

Montañez, I.P., Osleger, D.A., Banner, J.L., Mack, L., and Musgrove, M.L., 2000, Evolution of the Sr and C isotope composition of Cambrian oceans: *GSA Today*, v. 10, no. 5, p. 1–7.

Pearson, P.N., Ditchfield, P.W., Singano, J., Harcourt-Brown, K.G., Nicholas, C.J., Olsson, R.K., Shackleton, N.J., and Hall, M.A., 2001, Warm tropical sea surface temperatures in the Late Cretaceous and Eocene epochs: *Nature*, v. 413, p. 481–487.

Rahmstorf, S., Archer, D., Ebel, D.S., Eugster, O., Jouzel, J., Maraun, D., Neu, U., Schmidt, G.A., Severinghaus, J., Weaver, A.J., and Zachos, J.C., 2004, Cosmic rays, carbon dioxide, and climate: *Eos (Transactions, American Geophysical Union)*, v. 85, p. 38, 41.

Veizer, J., Ala, D., Azmy, K., Bruckschen, P., Buhl, D., Bruhn, F., Carden, G.A.F., Diener, A., Ebner, S., Godderis, Y., Jasper, T., Korte, C., Pawellek, F., Podlaha, O.G., and Strauss, H., 1999, ⁸⁷Sr/⁸⁶Sr, δ¹³C and δ¹⁸O evolution of Phanerozoic seawater: *Chemical Geology*, v. 161, p. 59–88.

Wenzel, B., 2000, Differential preservation of primary isotopic signatures in Silurian brachiopods from northern Europe: *Journal of Sedimentary Research*, v. 70, p. 194–209.